



Global glacier mass changes and their contributions to sea-level rise from 1961 to 2016

Zemp, M., Huss, M., Thibert, E., Eckert, N., McNabb, R., Huber, J., Barandun, M., Machguth, H., Nussbaumer, S. U., Gärtner-Roer, I., Thomson, L., Paul, F., Maussion, F., Kutuzov, S., & Cogley, J. G. (2019). *Global glacier mass changes and their contributions to sea-level rise from 1961 to 2016*. Abstract from European Geosciences Union General Assembly, Vienna, Austria.

[Link to publication record in Ulster University Research Portal](#)

Publication Status:

Published (in print/issue): 08/04/2019

Document Version

Publisher's PDF, also known as Version of record

General rights

Copyright for the publications made accessible via Ulster University's Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact pure-support@ulster.ac.uk.



Global glacier mass changes and their contributions to sea-level rise from 1961 to 2016

Michael Zemp (1), Matthias Huss (2,3), Emmanuel Thibert (4), Nicolas Eckert (4), Robert McNabb (5), Jacqueline Huber (1), Martina Barandun (3), Horst Machguth (1,3), Samuel Nussbaumer (1,3), Isabelle Gärtner-Roer (1), Laura Thomson (6), Frank Paul (1), Fabien Maussion (7), Stanislav Kutuzov (8), and Graham Cogley (9)

(1) University of Zurich, Department of Geography, Zurich, Switzerland (michael.zemp@geo.uzh.ch), (2) Laboratory of Hydraulics, Hydrology and Glaciology (VAW), ETH Zurich, Switzerland, (3) Department of Geosciences, University of Fribourg, Switzerland, (4) Université Grenoble Alpes, Irstea, UR ETGR, Grenoble, France, (5) Department of Geosciences, University of Oslo, Norway, (6) Department of Geography and Planning, Queen's University, Kingston, Canada, (7) Department of Atmospheric and Cryospheric Sciences, University of Innsbruck, Austria, (8) Department of Glaciology, Institute of Geography, Russian Academy of Sciences, Moscow, Russia, (9) Department of Geography, Trent University, Peterborough, Canada

Glaciers distinct from the Greenland and Antarctic Ice Sheets cover an area of approximately 706,000 km² globally with an estimated total volume of 170,000 km³, or 0.4 m of potential sea-level rise equivalent. Retreating and thinning glaciers are icons of climate change and impact regional runoff as well as global sea level. For the previous reports of the Intergovernmental Panel on Climate Change (IPCC), mass-change estimates were based on the multiplication of averaged or interpolated results from available observations of a few hundred glaciers with regional glacier areas. For data-scarce regions, these results had to be complemented with estimates based on satellite altimetry and gravimetry. These past approaches were challenged by the small number and heterogeneous spatio-temporal distribution of in situ measurement series and their often unknown representativeness for the respective mountain range as well as by spatial limitations of current satellite altimetry (only point data) and gravimetry (coarse resolution).

In view of the upcoming IPCC Special Report on the Ocean and Cryosphere in a Changing Climate (SROCC), we gathered an international team of experts to provide a new estimate of glacier contributions to global mean sea level. Here we present a new approach that combines the temporal variability from glaciological observations with improved spatial coverage from geodetic observations. Our new approach suggests that glaciers contributed 27 ± 22 mm to global mean sea-level rise from 1961 to 2016. Regional specific mass-change rates for 2006–2016 range between -0.1 and -1.2 m water equivalent per year, resulting in a global sea-level contribution of 335 ± 144 Gt per year or 0.92 ± 0.39 mm per year. The current glacier mass loss is thus equivalent to the sea-level contribution of the Greenland Ice Sheet, clearly exceeds the loss from the Antarctic Ice Sheet, and accounts for 25% to 30% of the total observed sea-level rise.